
Prepared in cooperation with the STATE OF NEW HAMPSHIRE Governor's Energy Office and Department of Environmental Services Dr. Eugene L. Boudette, State Geologist



UTM GRID AND 1985 MAGNETIC NORTH DECLINATION AT CENTER OF SHEET

EXPLANATION

Wetland Description

Peat labsent or generally less than 5 ft thick

Peat averages 5 ft thick

Peat averages between 5 and 10 ft thick

Peat averages more than 10 ft thick

¹The American Society for Testing and Materials (1969) defines commercial peat as having an ash content of not more than 25 per cent dry

Wetland number classified in table 1

Locality of core. See figure 1 for descriptions

PEAT RESOURCES, PEATLANDS AND WETLANDS

Concepts and Definitions

Peat is a light- to dark-brown or black residuum formed by the partial decay and disintegration of plants that grew in marshes and swamps or in other damp places, such as raised bogs commonly known as heaths in New England. Peat may be (1) fibrous, matted material composed of mosses, ferns, grasses, rushes, reeds, sedges, and woody material from trees and shrubs; (2) finely divided plant material so decomposed that their biological identity is lost; or (3) nonfibrous, plastic, colloidal, and macerated material deposited at the bottom of lakes or other bodies of

Resources:

Commercial quality peat is defined by The American Society for Testing Materials (ASTM) as containing an ash content of not more than 25 percent on an air-dried basis. This peat should be at least 5 feet thick to be considered as a possible resource.

Because peat is derived from different types of vegetation and may contain varying amounts of mineral matter, the properties and composition of peat can vary considerably in different deposits and even in different

parts of the same deposit. The principal factors that determine the commercial value of peat are: water-holding capacity, organic and ash content, fiber content, and acidity. The ASTM has published standard methods for testing each of these factors (ASTM-D-29) which may be obtained from the American Society for Testing Materials, 1916 Race St.,

Peat is mined throughout the world chiefly for use in agriculture and horticulture, and to a lesser extent as a fuel. Its value for manufactured goods and for use as a filter is under current investigation. Peatlands and Wetlands

Attention has recently focused on the value of in situ peat as an

important factor in the environmental control of wastes from mines and factories and contamination from agricultural herbicides and insecticides. Peat not only soaks up elements like a sponge, but chemical processes and organisms within the peat and in the peat-forming environment bring about changes in organic and chemical wastes drained into the deposits. Marshes, swamps, and bogs containing peat deposits of varying thicknesses are collectively known as peatlands. They are also grouped under the more generalized term of "wetlands" by Cowardin and others (1979). Wetlands are dscribed by these authors as lands where saturation with water is the dominant factor determining how soil is developed and what types of animal and plant communities live in the soil and on its surface. Wetlands are transitional between terrestrial and aquatic systems where the water table is usually at or near the surface; the land may be covered by shallow water at least seasonally. Wetlands may contain an organic-type soil so high in ash content that it is not considered peat.

Wetland/Peatland Classification Guide for New Hampshire

This guide is developed as an extension of the National Wetlands Classification System (Cowardin and others, 1979), which was designed by the U.S. Fish and Wildlife Service and is concerned primarily with wildlife habitats in which the wetland surface is of greatest importance. The three parameters of their classification is based are hydrophytic vegetation, hydric soil, and hydrology.

The extended classification used on this map adds the geologic parameters in a descriptive identification for both the wetland and its setting. The setting controls the stratigraphic development of the organic and inorganic material immediately below the wetland surface. The shape of the basin influences the thickness of both the organic and inorganic materials in the wetland. Type and structure of the surrounding consolidated and unconsolidated rock influence the ash and element content of the organic material beneath the wetland surface. Ground and surface water regimes also control ash and element content, as well as the stratigraphic development on which the amount of peat is predicted. Predictable chemical, bacterial, and physical processes are likewise basically controlled by shape and material of the rock basin and the ground and surface water regimes. The understanding of these processes, together with soil and vegetation controls, is vital to assessing the value of peatlands/wetlands for waste, herbicide, and pesticide control.

This extended classification system, like that of the Fish and Wildlife Service, is hierarchial. At the highest level "OTHER GEOLOGICAL ASPECTS" (see Table 1) have been added to their "HYDROPHIC VEGETATION," "HYDRIC SOIL," and "HYDROLOGY." Under each of these four major headings are appropriate modifiers and submodifiers (See example).

Example: Wetland 14, Core I

This wetland shown on the map and described in table 1 is located in Mount Vernon Township. It has marsh and swamp types of hydrophitic vegetation under which is a peat deposit of commercial quality; ash content is less than 25 percent on the dry basis. This wetland is subject to overflow, which means that silt is added from time to time. The vegetation cover changes as swamp trees are drowned when the water table rises and are replaced by marsh grasses following the draining of the site. The preglacial bedrock adjacent to the wetland is completely buried by glacial drift. Surficial geology is taken from Kotef (1979). The Ds1-6 bedrock symbol taken from Lyons and others (1986) stands for a weakly foliated to non-foliated spotted biotite quartz diorite, tonalite, granodiorite and granite with possible garnet and muscovite (Spaulding Quartz Diorite Intrusive Suite). The organic material of the wetland is peat covered by a marsh or swamp with a relatively flat surface, meaning that the more acidic and nutrient-poor conditions that give rise to raised moss- and heath-covered bogs have not developed since last flooding.

Table 2 shows that wetland 14 cored at I covers 45 acres with an average of 10 ft of commercial-quality peat. This layer amounts to a resource of an estimated 90,000 tons of air-dried peat. Thickness is facilitated by the steep walls of the basin which is filling up.

Tables 3 and 4 show analyses of eight samples from core I, the stratigraphy of which is illustrated in figure 1. Ash content generally increases with depth, meaning the original pond filled first with mineral matter washed into open water. Later, as the basin filled with organic matter, ash content decreased; acidity increases as shown by a decrease in pH. Sulfur content is 1 percent or below at all depths.

REFERENCES CITED

American Society for Testing Materials, 1969, D2607-69, Standard classification of peats, mosses, humus, and related products: 1916 Race St., Philadelphia, Pa. 19103, 1 p. Cowardin, L.M., Carter, Virginia, Golet, F.C., and LaRoe, E.T., 1979,

Classification of wetlands and deepwater habitats of the United States: U.S. Fish and Wildlife Service, p. 1-103. Lyons, J.B., Bothner, W.A., Moench, R.H., and Thompson, J.B., 1986,

Interim geologic map of New Hampshire: New Hampshire Department

of Environmental Services Map OS-1-86, scale 1:250,000. Koteff, Carl, 1970, Surficial geologic map of the Milford quadrangle, Hillsborough County, New Hampshire: U.S. Geological Survey Geologic Quadrangle Map GQ-881, scale 1:24,000.




Table 2.--Estimated peat resources [Resources are estimated on the basis of a minimum thickness of 5 fe and a 1-acre-foot yield of 200 tons air-dried peat]

									Dry basis						3	
					Proximate				Ultimate			Total S	Sulfur and S	ulfur for	ms	
Wetland	Core	Sample	Btu Per lb.	Ash %	Volitile %	Fixed Carbon	Carbon %	Hydrogen %	Nitrogen %	Chlorine %	Pyritic %	Sulfate %	Organic %	Total %	Moisture a received %	
14	I	87-80B	7750	24.27	52.06	23.67	41.54	4-15	2.18	0.4	0.01	0.13	0-44	0.58	88.96	
14	I	87-82B	6166	40.44	44.84	14.72	35.26	3.51	1.93	0.02	0.02	0.10	0.38	0.50	88.58	

Wetland Core Number letter	Core letter	I HYDROPHITIC VEGETA		ATION	II HYDRIC SOIL upper 5 ft		III HYDRO	water table	IV OTHER GEOLOGICAL ASPECTS																					
											A. Preglacial bedrock adjacent to wetland							B. Glacial deposits adjacent to wetlands					C. Post-glacial deposits					D. Topography of wetland		
	A	. Swamp	B.Marsh	C. Heath D. 7	D. Tidal	A. Peat with	B.Muck	A. Subject to	B. Not readily subject to overflow	C. Subject		1. Valley	1. Valley or basin wall slope		3. Structure		1. Slope			2. Texture		1.	1. Deposits adjacent to we		and	2. Wetland deposits				
		THE OWNING			Marsh	ash content (dry basis) less than 25%	(ash content greater	overflow including drowning by beaver or man		to pronounced water table shifts	a. Steep	b. Moderate	c. Gentle d. Complete	2. Type (symbol from New Hampshire Gelogic Map)	a. Many fractures	b. Along a fault trace	a. Gentle	b. Moderate	c. Steep	a. Coarse, e.g. sand, gravel		a. Stream	b. Dune	c. Tidal	d. Not applicable	a. Inorganic e.g sand, silt, clay	b. Organic (m) muck (p) peat	1. Flat e.g. 2. Rounded with heath surrounded swamp by moat	eath surrounded	3. Forest floo appearing flat a comparitively
1	L	Х				X	X	X		X	Х			De1m					Х	X					Х		(p)	Х		
2	M	X				X	X	X		х	Х			De1m					Х	X					Х		(p)	Х		
3	N	X	X			X	X	X		Х	Х			De1m		7		Х		X		Х					(p)	Х		
4	P	X	X			X	X	X		Х	X			Ds1-6					X	X	Х	Х					(p)	Х		
5	0			X		X			X				X	Ds1-6					Х	X	Х				Х		(p)		X	
6	G	X	X				Х	X		X			X	Ds1-6	H 3	Х		X		X	Х	Х					(m)	Х		
7	K	X	X				X	X		X	X			Ds1-6				X		Х		X					(m)	Х		
8	F	X				X			X				X	Ds1-6			Х			X					Х		(p)	Х		
9	Н	X					X	X					X	Ds1-6					Х	Х	Х				Х		(m)	X		
10	J			X		X			X		X			Ds1-6		Х			X	X	Х				Х		(p)		Х	
11	E	X				X			X		Х			Ds1-6				X		Х	Х				Х		(p)	X		
12	C&D	X	X				Х	X				Х		Ds1-6	I K I Z			Х		Х	Х	Х					(m)	X		
13	В	X	х				X						X	Ds1-6				Х		Х	X				Х		(m)	X		
14	I	X	X			X					X			Ds1-6					X	Х	Х	Х					(p)	X		
15	A	X	X			X							X	Ds1-6				X		Х	X				Х		(p)	Х		

QUADRANGLE LOCATION

[See map for locations of wetlands and cores. See figure 1 for locations of samples in cores] (total) 0.35 0.55 87-86 87-87 87-78 87-80 87-81 87-82 87-83 87-84

A 87-72 7.1 4.2

Table 4.--Ash, acidity, moisture, and sulfur content of core samples

WETLAND AND PEAT RESOURCE MAP OF THE NEW BOSTON 7.5-MINUTE QUADRANGLE, NEW HAMPSHIRE

By

Cornelia C. Cameron, J.I. Marlow, III, and David A. Emery